arXiv:1610.05277v1 [math.DG] 17 Oct 2016

SPACES OF HARMONIC MAPS OF THE PROJECTIVE PLANE TO THE FOUR-DIMENSIONAL SPHERE

RAVIL GABDURAKHMANOV

Abstract. This paper investigates the spaces of harmonic maps of the projective plane to the four-dimensional sphere by means of twistor lifts. It was shown that such spaces are empty in case of the even degree. In case of the degree less than 6 it was shown that such spaces are path-connected and it was found an explicit parameterization of the canonical representatives. In addition the last section summarizes known results for harmonic maps of the two-dimensional sphere to the four-dimensional sphere.

1. Introduction

Let $\phi : M \to N$ be a map between Riemannian manifolds. We define its *energy* by the formula

$$E(\phi) = \frac{1}{2}\int\limits_M |d\phi(x)|^2 dx,$$

where $d\phi(x)$ is the differential of ϕ at the point $x \in M$; and dx is the volume element of M. With the functional E associated Euler-Lagrange operator $\tau(\phi) = div(d\phi)$ called *tension field of* ϕ. The map $\phi : M \to N$ is said to be *harmonic* if its tension field vanishes identically so that ϕ is a critical point of E.

Harmonic maps come in a different ways in mathematics: in the theory of minimal surfaces, spectral geometry and etc. For example:

- If $dim\, M = 1$, then the harmonic maps are the geodesics of N.
- If $N = \mathbb{R}$, they are harmonic functions on M.
- If $dim\, M = 2$, they include parametric representations of the minimal surfaces of N; the energy is the Dirichlet-Douglas integral.

As a result there were many investigations in the theory of the harmonic maps. This paper based on the results of the famous works of E. Calabi [1, 2], J. Barbosa [3], R. L. Bryant [4], and essentially uses works of J. Bolton and L. M. Woodward [6, 7, 8]. We use following proposition and fundamental theorems.

Proposition 1.1. [9] *An isometric immersion $\phi : (M, g) \to (N, h)$ is minimal if and only if it is harmonic.*

Theorem 1.2. (*E. Calabi* [1]). *Let $X : S^2 \to \mathbb{R}^n$ be an immersion of a sphere S^2 into $\mathbb{R}^n$, whose image is a locally minimal surface in a sphere rS^{n-1} of radius r, and is not contained in any hyperplane of $\mathbb{R}^n$. Then the following conclusions hold:*

(1) The area $A = A(S^2)$ of the image sphere S^2 is an integer multiple of $2\pi r^2$.

(2) The dimension number n is odd.

In this theorem used assumption that image of $X : S^2 \to \mathbb{R}^n$ is not contained in any hyperplane of $\mathbb{R}^n$. Such maps are called *linearly full*. It should be noted that if we induce metric on S^2 by X in the conditions of theorem 1.2, then X is an isometric immersion, and hence harmonic due to 1.1. J. Barbosa clarifies paragraph (1) of theorem 1.2 in the case of $r = 1$.

Theorem 1.3. (*J. Barbosa* [3]). *The area of a generalized minimal immersion $X : S^2 \to S^{2m}$ is $4\pi d$, for some $d \in \mathbb{N}$.*

The word *generalized* in this theorem means that the metric, induced on S^2 by X, may be degenerated in isolated points. The number d, appearing in this theorem, is called *harmonic degree* (or *degree*) of harmonic map X. The case $m = 2$ has been investigated by R. L. Bryant in [4] and uses the construction of twistor fibration $\pi : \mathbb{C}P^3 \to S^4$ introduced in Section 2. The required results of his work could be summarized in the following theorem.

Theorem 1.4. (*R. L. Bryant* [4]).

(1) Let $\psi : S^2 \to \mathbb{C}P^3$ be a linearly full horizontal holomorphic curve then $\phi = \pi\psi : S^2 \to S^4$ is a linearly full harmonic map. Conversely every linearly full harmonic map $\phi : S^2 \to S^4$ is equal to $\pm\pi\psi$ for some uniquely determined linearly full horizontal holomorphic curve ψ called the twistor lift *of ϕ.*

(2) Twistor lift of linearly full harmonic map $\phi : S^2 \to S^4$ is a polynomial curve.

(3) Area of S^2 in the metric induced by ϕ is equal to $4\pi d$, and d is a degree of twistor lift.

This paper investigates harmonic maps $\phi : \mathbb{R}P^2 \to S^4$ by using standard Riemannian double cover $S^2 \to \mathbb{R}P^2$ and twistor lifts to $\mathbb{C}P^3$ of *antipodal-agreed* harmonic maps $S^2 \to S^4$ (i.e. harmonic maps that can be factorized through the double cover $S^2 \to \mathbb{R}P^2$).

2. The twistor fibration

We introduce the construction of the *twistor fibration* $\pi : \mathbb{C}P^3 \to S^4$. If $\mathbb{H}^2$ is a left quaternionic vector space, then π is obtained by composing the Hopf map $\rho : \mathbb{C}P^3 \to \mathbb{H}P^1$ given by

$$\rho([z_1, z_2, z_3, z_4]) = [z_1 + z_2 j, z_3 + z_4 j],$$

with the canonical identification of $\mathbb{H}P^1$ and $S^4 \subset \mathbb{H} \oplus \mathbb{R} = \mathbb{R}^5$ given by the stereographic projection from the south pole of S^4 onto equatorial hyperplane $\mathbb{H}$ in $\mathbb{R}^5$ which is included in $\mathbb{H}P^1$ by $q \mapsto [q, 1]$. This identification is given by

$$[q_1, q_2] \in \mathbb{H}P^1 \leftrightarrow \frac{(2\bar{q}_1 q_2, |q_1|^2 - |q_2|^2)}{|q_1|^2 + |q_2|^2} \in S^4.$$

Hence, we obtain an explicit formula for the twistor fibration

$$\pi([z_1, z_2, z_3, z_4]) = \frac{(2(\bar{z}_1 z_3 + z_2 \bar{z}_4), 2(\bar{z}_1 z_4 - z_2 \bar{z}_3), |z_1|^2 + |z_2|^2 - |z_3|^2 - |z_4|^2)}{|z_1|^2 + |z_2|^2 + |z_3|^2 + |z_4|^2}.$$

If $\mathbb{C}P^3$ is given the Fubini-Study metric of constant holomorphic sectional curvature 1 then π is a Riemannian submersion and the *horizontal distribution* on $\mathbb{C}P^3$ consists of those tangent vectors to $\mathbb{C}P^3$ which are orthogonal to the fibres of π.

Let

$$J = \begin{pmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \end{pmatrix}. \tag{2.1}$$

Then the projectivisation $PSp(2;\mathbb{C})$ of the complexified symplectic group

$$Sp(2;\mathbb{C}) = \{A \in SL(4;\mathbb{C}) | A^t J A = J\}$$

acts on $\mathbb{C}P^3$ in the standard way as the group of holomorphic diffeomorphisms, which preserve the horizontal distribution, with $PSp(2) = PSp(2;\mathbb{C}) \cap PU(4)$ the subgroup of holomorphic isometries which preserve the horizontal distribution [7].

It should be noted that all of these groups act transitively on $\mathbb{C}P^3$ and $Sp(2)$ and hence $PSp(2)$ are path-connected [10]. It should be noted also that the elements of $Sp(2)$ is matrices $(\mathbf{u}, J\bar{\mathbf{u}}, \mathbf{v}, J\bar{\mathbf{v}})$ such that $\mathbf{u}, \mathbf{v}$are unit vectors in $\mathbb{C}^4$, and $\mathbf{v}$ unitarily orthogonal to $\mathbf{u}$ and $J\bar{\mathbf{u}}$.

Let S be a Riemann surface and $\psi : S \to \mathbb{C}P^3$ holomorphic curve. Then ψ is *horizontal* if in each point it is tangential to the horizontal distribution on $\mathbb{C}P^3$, or equivalently if it intersects each fibre of π orthogonally. The condition [4] for a holomorphic curve $\psi = [\mathbf{f}] = [f_1, f_2, f_3, f_4]$ to be horizontal is

$$(\mathbf{f}', J\mathbf{f}) = 0, \tag{2.2}$$

where $(\ ,\ )$ denotes the complex bilinear extension to $\mathbb{C}^4$ of the standard real inner product on $\mathbb{R}^4$. The horizontality condition may be written as

$$f_1' f_2 - f_1 f_2' + f_3' f_4 - f_3 f_4' = 0. \tag{2.3}$$

Here and throughout the rest of the paper we write holomorphic curves $\psi : S \to \mathbb{C}P^n$ locally in terms of a local complex coordinate z in the form $\psi = [\mathbf{f}] =$

$[f_1, f_2, ..., f_{n+1}]$, with $f_1, ..., f_{n+1}$ holomorphic functions of z without common zeros. We will call such a representation a *reduced form* of ψ.

3. Higher singularities for horizontal holomorphic curves

We first recall the definition of singularity type of holomorphic curves in $\mathbb{C}P^n$ [6]. Suppose that $\psi : S \to \mathbb{C}P^n$ is a linearly full holomorphic curve. We write ψ locally in reduced form $\psi = [\mathbf{f}] = [f_1, f_2, ..., f_{n+1}]$, and let $\mathbf{f}^{(j)}$ denote the j-th derivative of $\mathbf{f}$ with respect to z. For each $k = 0, \ldots, n-1$ we define *k-th osculating curve* as $[\mathbf{f} \wedge \ldots \wedge \mathbf{f}^{(k)}]$. A *higher singularity* of ψ is a point $p \in S$ at which the derivative of k-th osculating curve is equal to zero for some $k = 0, \ldots, n-1$. The set $Z(\psi)$ of higher singularities of ψ is therefore given by

$$Z(\psi) = \{p \in S \,|\, \mathbf{f} \wedge \ldots \wedge \mathbf{f}^{(n)}(p) = 0\}.$$

If z is a local complex coordinate with $z(p) = 0$, then $\mathbf{f}$ can be written in the *normal form*

$$\mathbf{f}(z) = h_0(z)\mathbf{a}_0 + z^{r_0(p)+1}h_1(z)\mathbf{a}_1 + \ldots + z^{r_0(p)+\ldots r_{n-1}(p)+n}h_n(z)\mathbf{a}_n$$

for some suitable choise of basis $\mathbf{a}_0, \ldots, \mathbf{a}_n$ of $\mathbb{C}^{n+1}$, non-negative integers $r_0(p), \ldots, r_{n-1}(p)$, and holomorphic functions $h_0(z), \ldots, h_n(z)$, each non-zero at $z = 0$. If $r_k(p) > 0$ then the derivative of the k-th osculating curve has a zero of order $r_k(p)$ at p and the *singularity type* of ψ is defined to be the set

$$\{(p; r_0(p), \ldots, r_{n-1}(p)) \,|\, p \in Z(\psi)\}.$$

Lemma 3.1. [6] *Let $g \in PGL(n+1;\mathbb{C})$ and ω a conformal diffeomorphism of S. If ψ has singularity type $\{(p; r_0(p), \ldots, r_{n-1}(p)) \,|\, p \in Z(\psi)\}$ then $g\psi\omega^{-1}$ has singularity type $\{(\omega(p); r_0(p), \ldots, r_{n-1}(p)) \,|\, p \in Z(\psi)\}$.*

We now give a criterion for determining the higher singularities of a linearly full horizontal holomorphic curve $\psi : S \to \mathbb{C}P^3$.

Lemma 3.2. [6] *Let $\psi : S \to \mathbb{C}P^3$ be a linearly full horizontal holomorphic curve written locally in terms of a complex coordinate z in reduced form as $\psi(z) = [\mathbf{f}(z)] = [f_1(z), f_2(z), f_3(z), f_4(z)]$. Then $z = a$ is a higher singularity of ψ if and only if*

$$(f_1''f_2' - f_1'f_2'' + f_3''f_4' - f_3'f_4'')(a) = 0, \tag{3.1}$$

or equivalently, $(\mathbf{f}'', J\mathbf{f}')(a) = 0$.

For linearly full holomorphic curve $\psi : S^2 \to \mathbb{C}P^n$ and $k = 0, \ldots, n-1$ we define

$$r_k = \sum_{p \in Z(\psi)} r_k(p).$$

Proposition 3.3. *Let $\psi : S^2 \to \mathbb{C}P^3$ be a linearly full horizontal holomorphic curve and $p \in S^2$. Then $r_0(p) = r_2(p)$ and*

$$2r_0 + r_1 = 2d - 6, \tag{3.2}$$

where d is the degree of ψ.

Proofs of the lemmas and proposition of this section can be found in [6].

4. A canonical form

Due to 1.4 the twistor lift of linearly full harmonic map $\phi : S^2 \to S^4$, and in particular the fact that it is a polynomial curve, gives us a good opportunity for investigating harmonic maps. Namely, let $HHol_d^{LF}(\mathbb{C}P^3)$ denote the set of linearly full horizontal holomorphic maps of S^2 to $\mathbb{C}P^3$ of degree d, and let $Harm_d^{LF}(S^4)$ denote the set of linearly full harmonic maps of S^2 to S^4 with induced area $4\pi d$. Then $Harm_d^{LF}(S^4)$ is divided into two components $Harm_d^{+}(S^4)$ and $Harm_d^{-}(S^4)$, which can be interchanged by the antipodal involution of S^4, and there is a bijective correspondence

$$\pi_*^{\pm} : HHol_d^{LF}(\mathbb{C}P^3) \to Harm_d^{\pm}(S^4) \tag{4.1}$$

defined by $\pi_*^{\pm}(\psi) = \pm\pi \circ \psi$.

Clearly, $p \in S^2$ is a branch point of ψ if and only if $r_0(p) > 0$ while as is shown in Section 7 of [7], if $r_0(p) = 0$ then p is an umbilic of ψ if and only if $r_1(p) > 0$. Further, it follows from work [11] that the branch points and umbilics of ϕ coincide with those of ψ. Thus the higher singularities of ψ occur at the branch points or umbilics of ϕ.

It is clear that the group $PSp(2)$ acts freely on $HHol_d^{LF}(\mathbb{C}P^3)$ by postcomposition via its standard action on $\mathbb{C}P^3$ and it follows from lemma 3.1 that the singularity type is preserved. The identifications determined by $\pi_*^{\pm}$determine an action of $PSp(2)$ on $Harm_d^{\pm}(S^4)$ wich preserves induced area, branch points, umbilics and antipodal-agreement. Similarly the rotation group $SO(3) \cong PSU(2)$ also acts on $HHol_d^{LF}(\mathbb{C}P^3)$ and $Harm_d^{\pm}(S^4)$ by precomposition via its standard action on $S^2 = \mathbb{C}P^1$ and also preserves antipodal-agreement. Note that the actions of $PSp(2)$ and $PSU(2)$ commute and that the maps $\pi_*^{\pm}$ are $PSp(2)$-equivariant and $PSU(2)$-equivariant.

Now we consider the vector space $\mathbb{C}[z]_d$ of polynomials of degree at most d, and let V be the subset of $(\mathbb{C}[z]_d)^4$ consisting of those quadruples of coprime polynomials with maximum degree d for which the map $z \to [f_1(z), f_2(z), f_3(z), f_4(z)]$ is linearly full in $\mathbb{C}P^3$. Then V is a projective subset of $(\mathbb{C}[z]_d)^4$, and we identify its projectivisation $P(V)$ with the space of linearly full holomorphic maps of degree d

from S^2 to $\mathbb{C}P^3$ in the usual way via

$$[f_1, f_2, f_3, f_4] \leftrightarrow (z \to [f_1(z), f_2(z), f_3(z), f_4(z)]).$$

Here, and subsequently, we use the complex coordinate on S^2 defined by stereographic projection from the south pole of S^2 onto the equatorial plane so that, in the usual sense, we may identify S^2 with $\mathbb{C} \cup \{\infty\}$. Through this identification antipodal involution interchanges z and $-1/\bar{z}$.

We give $(\mathbb{C}[z]_d)^4$ its natural topology as a vector space, and $P((\mathbb{C}[z]_d)^4)$ the identification topology. Then V is open subset of $(\mathbb{C}[z]_d)^4$, and $P(V)$ is an open subset of $P((\mathbb{C}[z]_d)^4)$. Subsets of any of these spaces are then given the induced (subspace) topology. We denote by $V^H \subset V$ the subset of horizontal maps. Next we give $Harm_d^{LF}(S^4)$ the compact-open topology and rewrite lemma from [8].

Lemma 4.1. *$Harm_d^{\pm}(S^4)$ is a closed subsets of $Harm_d^{LF}(S^4)$, and the maps $\pi_*^{\pm}$: $P(V^H) \to Harm_d^{\pm}(S^4)$ are homeomorphisms.*

For convenience, we introduce *coefficient matrix (*or *matrix)* of the holomorphic curve $\mathbf{f}(z) = (f_1(z), f_2(z), f_3(z), f_4(z)) \in (\mathbb{C}[z]_d)^4$. It is given by

$$F = \begin{pmatrix} a_1^0 & a_1^1 & \dots & a_1^{d-1} & a_1^d \\ a_2^0 & a_2^1 & \dots & a_2^{d-1} & a_2^d \\ a_3^0 & a_3^1 & \dots & a_3^{d-1} & a_3^d \\ a_4^0 & a_4^1 & \dots & a_4^{d-1} & a_4^d \end{pmatrix},$$

where a_i^j is a j-th coefficient of the polynom $f_i(z) = a_i^0 + a_i^1 z + \ldots + a_i^j z^j + \ldots + a_i^d z^d$, and $1 \le i \le 4$, $0 \le j \le d$. From definition it is clear that $(\mathbf{f}(z))^t = F \cdot (1, z, z^2, \ldots, z^d)^t$ and also that $\mathbf{f}(z)$ is linearly full if the coefficient matrix F has a full rank.

Note that element $A \in Sp(2)$, acting on $\mathbf{f}(z) \in (\mathbb{C}[z]_d)^4$ by postcomposition (i.e. $A : \mathbf{f}(z) \to A \circ \mathbf{f}(z)$), acts on coefficient matrix F by the left multiplication (i.e. $A : F \to AF$). Also, as noted earlier, the group $PSU(2) \cong SO(3)$ acts on S^2 as a group of rotations. This action defines an action of $PSU(2)$ on $\mathbb{R}P^2$ since it preserves the antipodal points. It should be noted that the group $PSU(2)$ is path-connected[10].

In the following sections we will investigate linearly full harmonic maps of S^2 to S^4 of degrees $d = 3$, 4, 5 and 6, which are the liftings of harmonic maps of $\mathbb{R}P^2$ to S^4 through the double cover (i.e. antipodal-agreed maps). Let $AHarm_d^{LF}(S^4) \subset Harm_d^{LF}(S^4)$ denote the space of antipodal-agreed linearly full harmonic maps of S^2 to $S^4$of degree d, with $AHarm_d^{\pm}(S^4) \subset Harm_d^{\pm}(S^4)$. And let denote the twistor lifts of this spaces as $AHHol_d^{LF}(\mathbb{C}P^3) \subset HHol_d^{LF}(\mathbb{C}P^3)$. Our main idea is in deformation of elements of $AHarm_d^{\pm}(S^4) \subset Harm_d^{\pm}(S^4)$ to some canonical form

by the action of some appropriate elements of $PSU(2)$ and $PSp(2)$. Note that this groups act continuously on $AHHol_d^{LF}(\mathbb{C}P^3)$ and, due to their path-connectedness, we can recover some information about path-connectedness of $AHHol_d^{LF}(\mathbb{C}P^3)$.

5. Harmonic maps of arbitrary degree

In this section we will exclude the condition of antipodal-agreement by finding the general form of antipodal-agreed maps. In this way we will need some addition of the theorem 1.4.

Theorem 5.1. (*R. L. Bryant* [4]).

Let $\hat{\psi} : S^2 \to \mathbb{C}P^3$ be a linearly full horizontal antiholomorphic curve then $\phi = \pi\hat{\psi} : S^2 \to S^4$ is a linearly full harmonic map. Conversely every linearly full harmonic map $\phi : S^2 \to S^4$ is equal to $\pm\pi\hat{\psi}$ for some uniquely determined linearly full horizontal antiholomorphic curve $\hat{\psi}$ called the antiholomorphic twistor lift *of ϕ.*

Moreover if $\psi = [\mathbf{f}] = [f_1, f_2, f_3, f_4]$ is a twistor lift of ϕ then $\hat{\psi} = \bar{\psi}J = [\bar{\mathbf{f}}J] = [\bar{f}_2, -\bar{f}_1, \bar{f}_4, -\bar{f}_3]$ is an antiholomorphic twistor lift of ϕ [7].

For $\mathbf{f}(z) = (f_1(z), f_2(z), f_3(z), f_4(z))$ of degree n we introduce

$$\tilde{\mathbf{f}}(z) = (\tilde{f}_1(z), \tilde{f}_2(z), \tilde{f}_3(z), \tilde{f}_4(z)) = \\ = (\bar{z}^n f_1(-1/\bar{z}), \bar{z}^n f_2(-1/\bar{z}), \bar{z}^n f_3(-1/\bar{z}), \bar{z}^n f_4(-1/\bar{z})).$$

Then $[\tilde{f}(z)]$ is an antiholomorphic curve in $\mathbb{C}P^3$.

Theorem 5.2. *There is no harmonic maps of real projective plane to the four-dimensional sphere of even degree, i.e. spaces $AHarm_n^{LF}(S^4)$ are empty for even n.*

Proof. Let us suppose that such maps exist. Then for each map $\phi \in AHarm_n^{LF}(S^4)$ we will compose its twistor lift $\psi = [\mathbf{f}(z)] = [f_1(z), f_2(z), f_3(z), f_4(z)]$ with antipodal involution of S^2 and get an antiholomorphic twistor lift $\tilde{\psi} = [\tilde{\mathbf{f}}(z)]$. Using theorem 5.1 we get $\psi = -\bar{\tilde{\psi}}J$. Then

$$g(z) \cdot \mathbf{f}(z) = g(z) \cdot (f_1(z), f_2(z), f_3(z), f_4(z)) = \\ = -\overline{\tilde{\mathbf{f}}(z)}J = (-\overline{\tilde{f}_2(z)}, \overline{\tilde{f}_1(z)}, -\overline{\tilde{f}_4(z)}, \overline{\tilde{f}_3(z)}), \tag{5.1}$$

for some function $g(z)$ on S^2. Now we will prove that $g(z)$ is a holomorphic function. From (5.1) we have

$$g(z) = \frac{-\overline{\tilde{f}_2(z)}}{f_1(z)} = \frac{\overline{\tilde{f}_1(z)}}{f_2(z)} = \frac{-\overline{\tilde{f}_4(z)}}{f_3(z)} = \frac{\overline{\tilde{f}_3(z)}}{f_4(z)}. \tag{5.2}$$

It is clear that $\overline{\tilde{f}_i(z)}$ $(i = 1, 2, 3, 4)$ are polynomials of z of degree less or equal to n, so they are holomorphic functions on $\mathbb{C}$. At the same time $f_i(z)$ $(i = 1, 2, 3, 4)$ don't

have common roots and therefore $g(z)$ is holomorphic on $\mathbb{C}$. By the definition there is polynomial of degree n among $f_i(z)$ $(i = 1, 2, 3, 4)$ and from (5.2) we conclude that $g(z)$ is holomorphic at ∞. Thus $g(z)$ is the holomorphic function on S^2 and therefore $g(z) = \frac{1}{\beta} = const \neq 0$.

Since $f_i(z) = a_i^0 + a_i^1 z + \ldots + a_i^j z^j + \ldots + a_i^n z^n$, $(i = 1, 2, 3, 4)$ we have $\overline{\tilde{f}_i(z)} = (-1)^n \overline{a_i^n} + (-1)^{n-1} \overline{a_i^{n-1}} z + \ldots + (-1)^{n-j} \overline{a_i^{n-j}} z^j + \ldots + \overline{a_i^0} z^n$, $(i = 1, 2, 3, 4)$. Considering (5.2) we can write

$$\begin{cases} a_1^n = -\beta \overline{a_2^0}, & a_2^0 = (-1)^n \beta \overline{a_1^n}, \\ a_2^n = \beta \overline{a_1^0}, & a_1^0 = (-1)^{n+1} \beta \overline{a_2^n}, \\ a_3^n = -\beta \overline{a_4^0}, & a_4^0 = (-1)^n \beta \overline{a_3^n}, \\ a_4^n = \beta \overline{a_3^0}, & a_3^0 = (-1)^{n+1} \beta \overline{a_4^n}. \end{cases} \tag{5.3}$$

From these relations we can conclude for leading coefficients $a_i^n = (-1)^{n+1} |\beta|^2 a_i^n$ and since n is even $a_i^n = -|\beta|^2 a_i^n$. But it can only be when $a_i^n = 0$ $(i = 1, 2, 3, 4)$ and thus the degree of the curve is reducing. So we have the contradiction which proves this theorem. □

Remark 5.3. In case of odd n for leading coefficients we have $a_i^n = |\beta|^2 a_i^n$, therefore $|\beta| = 1$ and $\beta = \exp(i\tilde{\beta})$ for some $\tilde{\beta} \in \mathbb{R}$. In this way any curve of odd degree is of the form $\psi = [\mathbf{f}(z)] = [f_1(z), \beta \overline{\tilde{f}_1(z)}, f_3(z), \beta \overline{\tilde{f}_3(z)}]$. Note that if we act on the curve by rotation $\omega(z) = \exp(-\mathrm{i}\frac{\tilde{\beta}}{\mathrm{n}})z$, we get the curve with $\beta = 1$. (Note also that this rotation leaves points 0 and ∞ on their places). Thus any curve of odd degree can be rotated to the curve of form $\psi = [\mathbf{f}(z)] = [f_1(z), \overline{\tilde{f}_1(z)}, f_3(z), \overline{\tilde{f}_3(z)}]$.

In next sections we will need following usefull lemma.

Lemma 5.4. *Any curve $\psi \in HHol_d^{LF}(\mathbb{C}P^3)$ by appropriate element $g \in PSp(2) \subset PSp(2, \mathbb{C})$ can be reduced to the curve $g\psi$ with coefficient matrix of form*

$$\begin{pmatrix} * & * & * & * & \ldots & * & * & * & * \\ 0 & 0 & 0 & * & \ldots & * & * & * & * \\ 0 & * & * & * & \ldots & * & * & * & * \\ 0 & 0 & * & * & \ldots & * & * & * & * \end{pmatrix}.$$

In addition, if $\psi \in AHHol_d^{LF}(\mathbb{C}P^3) \subset HHol_d^{LF}(\mathbb{C}P^3)$ then by appropriate element $g \in PSp(2)$ it can be reduced to the curve $g\psi$ with coefficient matrix of form

$$\begin{pmatrix} * & * & * & * & \ldots & * & 0 & 0 & 0 \\ 0 & 0 & 0 & * & \ldots & * & * & * & * \\ 0 & * & * & * & \ldots & * & * & 0 & 0 \\ 0 & 0 & * & * & \ldots & * & * & * & 0 \end{pmatrix}.$$

Proof. First, transitivity of action of $PSp(2)$ on $\mathbb{C}P^3$ allows us to turn the first column to the needed form. Next we notice that the elements of $Sp(2)$ which preserve vector $(1,0,0,0)^t$ have a form

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & v_1 & -\bar{v}_2 \\ 0 & 0 & v_2 & \bar{v}_1 \end{pmatrix},$$

and since we have matrix from $U(2)$ in the right bottom corner, we can eliminate the last term of the second column. Other two zeros in the second row is due to the horizontality condition (2.3). In the addition zeros in the right side of the coefficient matrix is due to remark 5.3. □

6. Spaces of linearly full harmonic maps of degree less than 6

In previous sections we have shown that there aren't linearly full harmonic maps of $\mathbb{R}P^2$ to S^4 of even degree. Also it is clear that there aren't linearly full maps of degree less than 3. So we need to describe the spaces of such maps of degree 3 and 5. Here we have the following results.

Theorem 6.1. *Every linearly full horizontal holomorphic curve $\psi \in AHHol_3^{LF}(\mathbb{C}P^3)$ by the action of appropriate elements $g \in PSp(2)$ and $\omega \in PSU(2)$ can be reduced to the canonical form $\Psi^3(z) = g\psi\omega(z) = [1, -z^3, \sqrt{3}z, \sqrt{3}z^2]$.*

Proof. Using lemma 5.4 and remark 5.3 we get the coefficient matrix

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & a & 0 & 0 \\ 0 & 0 & -\bar{a} & 0 \end{pmatrix}.$$

Then, acting on this curve with $\omega(z) = -z$, $g = diag\{1, 1, \exp(-i\mathrm{Arg(a)}), \exp(i\mathrm{Arg(a)})\}$, we get

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & -a & 0 & 0 \\ 0 & 0 & -a & 0 \end{pmatrix},$$

with $a \in \mathbb{R}$. Checking horizontality condition (2.3) we conclude $a^2 = 3$. Thus we have two cases

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & \sqrt{3} & 0 & 0 \\ 0 & 0 & \sqrt{3} & 0 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & -\sqrt{3} & 0 & 0 \\ 0 & 0 & -\sqrt{3} & 0 \end{pmatrix}$$

which can be interchanged by $g = diag\{1, 1, -1, -1\}$. This completes the proof. □

Corollary 6.2. *Spaces $AHHol_3^{LF}(\mathbb{C}P^3)$ and $AHarm_3^{\pm}(S^4)$ are path-connected.*

Proof. Follows from theorem 6.1, lemma 4.1 and path-connectedness and continuity of actions of $PSp(2)$ and $PSU(2)$. □

So, the spaces of linearly full harmonic maps of $\mathbb{R}P^2$ to S^4 of area 6π in the induced metric have two path-connected components $AHarm_3^+(S^4)$ and $AHarm_3^-(S^4)$, which are homeomorphic through the antipodal involution of S^4.

Theorem 6.3. *Every linearly full horizontal holomorphic curve $\psi \in AHHol_5^{LF}(\mathbb{C}P^3)$ by the action of appropriate elements $g \in PSp(2)$ and $\omega \in PSU(2)$ can be reduced to the canonical form $\Psi_\eta^5(z) = g\psi\omega(z) = [1 + qz, -qz^4 + z^5, \mu z + \eta z^2, \eta z^3 - \mu z^4]$, where $q = \frac{\eta}{\sqrt{3}}\sqrt{\frac{\eta^2+5}{\eta^2+4}}$, $\mu = \frac{2}{\sqrt{3}}\sqrt{\frac{\eta^2+5}{\eta^2+4}}$, $\eta \in (-\infty, +\infty)$.*

Proof. First, we can conclude from equation (3.2) that such curves have four higher singularities. Certainly, there are two pairs of antipodal points. We choose one of these points and rotate it to 0 by appropriate element of $PSU(2)$. Next using lemma 5.4 and remark 5.3 we get for the coefficient matrix

$$\begin{pmatrix} 1 & q & p & 0 & 0 & 0 \\ 0 & 0 & 0 & \bar{p} & -\bar{q} & 1 \\ 0 & \mu & \eta & \nu & 0 & 0 \\ 0 & 0 & -\bar{\nu} & \bar{\eta} & -\bar{\mu} & 0 \end{pmatrix}.$$

Then writing horizontality condition (2.3) we get system of equations

$$\begin{cases} -3p + \nu\bar{\mu} = 0, \\ 2q - p\bar{q} - \eta\bar{\mu} = 0, \\ -3q\bar{q} + p\bar{p} + 5 - 3\mu\bar{\mu} + \eta\bar{\eta} + \nu\bar{\nu} = 0, \end{cases} \tag{6.1}$$

Using equation (3.1) and taking into account (6.1) we get the equation for higher singularities

$$pz^4 + 2qz^3 + (5 - q\bar{q} - \mu\bar{\mu})z^2 - 2\bar{q}z + \frac{\mu\bar{\nu}}{3} = 0. \tag{6.2}$$

Since we have higher singularity at 0, we get $\mu\bar{\nu} = 0$. Then from the first equation of (6.1) follows $p = 0$. Let us suppose $\mu = 0$, then from the second equation of (6.1) we get $q = 0$, but this contradicts with last equation of (6.1). Thus we have $\nu = 0$.

The next step is in turning non-zero coefficients into the reals. Let $q = \tilde{q}e^{ix}$, then using multiplication by $e^{i5/2x}$ and actions of $\omega(z) = \exp(-ix)z$, $g = diag\{e^{-5/2x}, e^{5/2x}, 1, 1\}$ we get $q \in \mathbb{R}$. Further, we use the action of $g = diag\{1, 1, e^{-i\mathrm{Arg}(\mu)}, e^{i\mathrm{Arg}(\mu)}\}$ and get $\mu \in \mathbb{R}$. We conclude from the second equation of (6.1) that $\eta \in \mathbb{R}$ also. So we have the coefficient matrix

$$\begin{pmatrix} 1 & q & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -q & 1 \\ 0 & \mu & \eta & 0 & 0 & 0 \\ 0 & 0 & 0 & \eta & -\mu & 0 \end{pmatrix}$$

and equations

$$\begin{cases} 2q - \eta\mu = 0, \\ -3q^2 + 5 - 3\mu^2 + \eta^2 = 0, \end{cases} \tag{6.3}$$

with real coefficients. The solutions of these equations are $\mu = \pm\frac{2}{\sqrt{3}}\sqrt{\frac{\eta^2+5}{\eta^2+4}}$, $q = \pm\frac{\eta}{\sqrt{3}}\sqrt{\frac{\eta^2+5}{\eta^2+4}}$, $\eta \in \mathbb{R}$. This pair of families of the solutions can be interchanged by multiplication on i and actions of $\omega(z) = -z$, $g = diag\{i, -i, i, -i\}$. This completes the proof. □

Corollary 6.4. *Spaces $AHHol_5^{LF}(\mathbb{C}P^3)$ and $AHarm_5^{\pm}(S^4)$ are path-connected.*

Proof. We can see that canonical form $\Psi_\eta^5(z)$ is continously dependent on η thus there is a path connecting any canonical form with $\Psi_{\eta=0}^5(z)$. With this fact our statement follows from theorem 6.3, lemma 4.1 and path-connectedness and continuity of actions of $PSp(2)$ and $PSU(2)$. □

So the spaces of linearly full harmonic maps of $\mathbb{R}P^2$ to S^4 of area 10π in the induced metric have two path-connected components $AHarm_5^+(S^4)$ and $AHarm_5^-(S^4)$, which are homeomorphic through the antipodal involution of S^4.

Remark 6.5. Under the conditions of theorem 6.3 the coefficient matrix of the canonical form for $\eta = 0$ is

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & -\sqrt{5/3} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \sqrt{5/3} & 0 \end{pmatrix},$$

and herewith four umbilic points with $r_1(p) = 1$ glued in two ombilic points with $r_1(0) = 2$, $r_1(\infty) = 2$.

On the other hand if we put $\eta \to +\infty$ then degree of the curve is reducing to 3 and we have matrix

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & \sqrt{3} & 0 & 0 \\ 0 & 0 & \sqrt{3} & 0 \end{pmatrix},$$

which is the matrix of canonical curve of degree 3 from theorem 6.1.There we have annihilation of pairs of umbilic points on 0 and ∞. This fact brings us to the following

Theorem 6.6. *The space $AHHol^{LF}_{\leq 5}(\mathbb{C}P^3)$ is path-connected.*

Proof. Follows from theorem 5.2, corollaries 6.2, 6.4 and remark 6.5. Indeed, spaces $AHHol^{LF}_{\leq 2}(\mathbb{C}P^3)$ and $AHHol^{LF}_{4}(\mathbb{C}P^3)$ are empty, but spaces $AHHol^{LF}_{5}(\mathbb{C}P^3)$ and $AHHol^{LF}_{3}(\mathbb{C}P^3)$ can be connected, for example, by the path

$$\gamma(t) = \begin{cases} \Psi^5_{tg(\pi t/2)}(z), & 0 \leq t < 1, \\ \Psi^3(z), & t = 1. \end{cases}$$

which will be continuous by remark 6.5. □

7. Linearly full harmonic maps of the two-dimensional sphere to the four-dimensional sphere of degree less than 6.

In this section we summarize known results for linearly full harmonic maps $S^2 \to S^4$ of degree less than 6. In this case we have groups $PSp(2,\mathbb{C})$ and $PSL(2,\mathbb{C})$ acting on the spaces $HHol^{LF}_{d}(\mathbb{C}P^3)$ by postcomposition and precomposition respectively. Note that these groups are path-connected [10]. For the beginning we provide usefull lemma from [6].

Lemma 7.1. [6] *The group $PSp(2,\mathbb{C})$ acts transitively on $\mathbb{C}P^3$ and the subgroup which fixes the point $[1,0,0,0]^t$ consists of the matrices of the form*

$$\begin{pmatrix} \xi & \eta & \xi(\gamma\lambda - \alpha\mu) & \xi(\delta\lambda - \beta\mu) \\ 0 & \zeta & 0 & 0 \\ 0 & \lambda & \alpha & \beta \\ 0 & \mu & \gamma & \delta \end{pmatrix}, \quad \alpha\delta - \beta\gamma = \xi\zeta = 1.$$

It is clear that there are no harmonic maps of degree less than 3. So we start with degree 3. Due to equation (3.2) we deduce that such maps can not have higher singularities. In this simple case we have following result.

Theorem 7.2. *Every linearly full horizontal holomorphic curve $\psi \in HHol^{LF}_{3}(\mathbb{C}P^3)$ by the action of appropriate elements $g \in PSp(2,\mathbb{C})$ and $\omega \in PSL(2,\mathbb{C})$ can be reduced to the canonical form $\tilde{\Psi}^3(z) = g\psi\omega(z) = [1, -z^3, \sqrt{3}z, \sqrt{3}z^2]$.*

Proof. Using lemma 5.4 we get coefficient matrix of form

$$\begin{pmatrix} 1 & a & b & c \\ 0 & 0 & 0 & d \\ 0 & r & p & q \\ 0 & 0 & s & t \end{pmatrix}. \tag{7.1}$$

Next, we use lemma 7.1 multiple times. After the each time, we rename all non-zero coefficients as they were before. So our sequence of transformation matrices from lemma 7.1 is following

$$\begin{pmatrix} 1 & 0 & {}^{t}\!/_{d} & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & d^{-1} & 0 \\ 0 & -t & 0 & d \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 & -{}^{q}\!/_{d} \\ 0 & 1 & 0 & 0 \\ 0 & -q & d & 0 \\ 0 & 0 & 0 & d^{-1} \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & s & -p \\ 0 & 0 & 0 & s^{-1} \end{pmatrix}, \\ \begin{pmatrix} d & -c & 0 & 0 \\ 0 & d^{-1} & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} \sqrt{-d} & 0 & 0 & 0 \\ 0 & \sqrt{-d}^{-1} & 0 & 0 \\ 0 & 0 & \sqrt{\frac{s}{r}} & 0 \\ 0 & 0 & 0 & \sqrt{\frac{r}{s}} \end{pmatrix}. \tag{7.2}$$

This sequence of transformations gives us following sequence of coefficient matrices

$$\begin{pmatrix} 1 & a & b & c \\ 0 & 0 & 0 & d \\ 0 & r & p & 0 \\ 0 & 0 & s & t \end{pmatrix} \to \begin{pmatrix} 1 & a & b & c \\ 0 & 0 & 0 & d \\ 0 & r & p & 0 \\ 0 & 0 & s & 0 \end{pmatrix} \to \begin{pmatrix} 1 & a & b & c \\ 0 & 0 & 0 & d \\ 0 & r & 0 & 0 \\ 0 & 0 & s & 0 \end{pmatrix} \to \\ \to \begin{pmatrix} 1 & a & b & 0 \\ 0 & 0 & 0 & d \\ 0 & r & 0 & 0 \\ 0 & 0 & s & 0 \end{pmatrix} \to \begin{pmatrix} 1 & a & b & 0 \\ 0 & 0 & 0 & -1 \\ 0 & s & 0 & 0 \\ 0 & 0 & s & 0 \end{pmatrix}. \tag{7.3}$$

Now we use the horizontality condition equation (2.3) and conclude that $a = b = 0$, $\frac{s^2}{3} = 1$. We have $s = \pm\sqrt{3}$ and these cases can be interchanged by $g = diag\{1, 1, -1, -1\}$. This completes the proof. □

As for the projective plane we have the following.

Corollary 7.3. *Spaces $HHol_3^{LF}(\mathbb{C}P^3)$ and $Harm_3^{\pm}(S^4)$ are path-connected.*

Proof. Follows from theorem 7.2, lemma 4.1 and path-connectedness and continuity of actions of $PSp(2, \mathbb{C})$ and $PSL(2, \mathbb{C})$. □

Here we see that canonical form $\tilde{\Psi}^3(z)$ for S^2 is equal to canonical form $\Psi^3(z)$ for $\mathbb{R}P^2$.

We continue with the case of degree 4. From equation (3.2) we conclude that such maps can have 1 or 2 higher singularities. We will see later that case of one higher singularity is impossible. In fact we have following.

Theorem 7.4. *Every linearly full horizontal holomorphic curve $\psi \in HHol_4^{LF}(\mathbb{C}P^3)$ by the action of appropriate elements $g \in PSp(2,\mathbb{C})$ and $\omega \in PSL(2,\mathbb{C})$ can be reduced to the canonical form $\tilde{\Psi}_a^4(z) = g\psi\omega(z) = [1{+}az, -z^4, \sqrt{2}z(1{+}{}^3\!/_2 az), \sqrt{2}z^3]$, $a \in \mathbb{R}$.*

Proof. For the beginning we turn one of the higher singularities to zero by appropriate element $\omega \in PSU(2)$. Next we use lemma 5.4 and get coefficient matrix of form

$$\begin{pmatrix} 1 & a & b & e & c \\ 0 & 0 & 0 & f & d \\ 0 & r & m & p & q \\ 0 & 0 & l & s & t \end{pmatrix}. \tag{7.4}$$

From equation (3.1) and because of higher singularity at zero, we conclude that $rl = 0$. It is clear that without loss of generality we may assume that $l = 0$. Horizontality condition equation (2.3) gives us $f = -\frac{rl}{3} = 0$. So, in fact our coefficient matrix is of form

$$\begin{pmatrix} 1 & a & b & e & c \\ 0 & 0 & 0 & 0 & d \\ 0 & r & m & p & q \\ 0 & 0 & 0 & s & t \end{pmatrix}. \tag{7.5}$$

Next we use sequence of transformations equation (7.2) and get coefficient matrix

$$\begin{pmatrix} 1 & a & b & e & 0 \\ 0 & 0 & 0 & 0 & -1 \\ 0 & s & m & 0 & 0 \\ 0 & 0 & 0 & s & 0 \end{pmatrix}. \tag{7.6}$$

There horizontality condition equation (2.3) gives us $b = e = 0$, $3a = ms$, $s^2 = 2$. So $s = \pm\sqrt{2}$ and these two cases can be interchanged by $g = diag\{1, 1, -1, -1\}$. We choose $s = \sqrt{2}$, then $m = {}^3\!/_{\sqrt{2}}a$. Now let $x = \mathrm{Arg}(a)$, then using multiplication by e^{i2x} and actions of $g = \{e^{-i2x}, e^{i2x}, e^{-ix}, e^{ix}\}$, $\omega(z) = e^{-ix}z$ we get $a \in \mathbb{R}$. This completes the proof. □

Remark 7.5. In this theorem we see that canonical form $\tilde{\Psi}_a^4(z)$ is continously dependent on a, therefore we can deform each one to the form $\tilde{\Psi}_0^4(z) = [1, -z^4, \sqrt{2}z, \sqrt{2}z^3]$. The equation equation (3.1) of higher singularities for these curves is $z(az+2) = 0$,

therefore second singularity is at point $z_0 = -2/a$. Note that second singularity for $\tilde{\Psi}_0^4(z)$ is at infinity. On the other hand, if we want to turn second singularity to zero, we must turn a to infinity, but then degree of the curve is decreasing to 1. So we conclude that such curves can not have one higher singularity. Therefore, in fact, we can deform every curve of degree 4 to $\tilde{\Psi}_0^4(z)$ by some element $g \in PSp(2,\mathbb{C})$ and Moebius transformation $\omega \in PSL(2,\mathbb{C})$ which turns higher singularities to the points 0 and ∞.

This remark immediately leads us to the following.

Corollary 7.6. *Spaces $HHol_4^{LF}(\mathbb{C}P^3)$ and $Harm_4^{\pm}(S^4)$ are path-connected.*

Proof. Follows from theorem 7.4, lemma 4.1, remark 7.5 and path-connectedness and continuity of actions of $PSp(2,\mathbb{C})$ and $PSL(2,\mathbb{C})$. □

Note that canonical forms for cases of dergees 3 and 4 coinside with the following theorem from [6].

Theorem 7.7. [6] *Let $\psi : S^2 \to \mathbb{C}P^3$ be a linearly full horizontal holomorphic curve of degree d which has at most two higher singularities. Then there exist elements $g \in PSp(2,\mathbb{C})$ and $\omega \in PSL(2,\mathbb{C})$ such that $\Psi(z) = g\psi\mu(z) = [1, k_2 z^{2k_1+k_2}, -(2k_1 + k_2)z^{k_1}, z^{k_1+k_2}]$, for some positive integers k_1, k_2 with $2k_1 + k_2 = d$. Higher singularities of $\Psi(z)$, if any, occur at points 0 and ∞.*

We start description of maps of degree 5 with the simple case. In this case we consider ψ has at most two higher singularities. Here we have following result.

Theorem 7.8. *Let $\psi : S^2 \to \mathbb{C}P^3$ be a linearly full horizontal holomorphic curve of degree 5 which has at most two higher singularities. Then there exist elements $g \in PSp(2,\mathbb{C})$ and $\omega \in PSL(2,\mathbb{C})$ reducing ψ to the one of the following form*

$$\tilde{\Psi}_1^5(z) = [1, z^5, 2z^2, -\frac{5}{2}z^3], \tag{7.7}$$

$$\tilde{\Psi}_2^5(z) = [1, z^5, z, -\frac{5}{3}z^4]. \tag{7.8}$$

Proof. From theorem 7.7 we get two cases $\Psi(z) = [1, z^5, -5z^2, z^3]$ and $\Psi(z) = [1, 3z^5, -5z, z^4]$ which can be reduced to (7.7) and (7.8) by elements $g_1 = diag\{1, 1, -\frac{2}{5}, -\frac{5}{2}\}$ and $g_2 = diag\{\sqrt{3}, \frac{1}{\sqrt{3}}, -\frac{\sqrt{3}}{5}, -\frac{5}{\sqrt{3}}\}$ respectively. □

It is worth mention that, in fact, both of forms from this theorem have two higher singularities.

So, we can consider the case of at least two higher singularities. In this most general case we have following result.

Theorem 7.9. [6] *Let $\psi : S^2 \to \mathbb{C}P^3$ be a linearly full horizontal holomorphic curve of degree 5. Then there exist elements $g \in PSp(2,\mathbb{C})$ and $\omega \in PSL(2,\mathbb{C})$ such that*

$$\tilde{\Psi}^5(z) = g\psi\mu(z) = [1 + az, (h+z)z^4, (r+lz)z, (m+sz)z^3] \tag{7.9}$$

canonical linearly full horizontal holomorphic curve, where a, h, r, l, m, s satisfy

$$2a + ls = 0,\ 5 + 3ah + lm + 3rs = 0,\ 2h + rm = 0. \tag{7.10}$$

Remark 7.10. Degree of curve from parametric family {(7.9), (7.10)} decreases iff $ah = 1$, $rs = lm = -2$, $hl = r$ or, equvivalently, $ah = 1$, $rs = lm = -2$, $ar = l$.

Lemma 7.11. *Every canonical form $\tilde{\Psi}^5(z)$ from theorem 7.9 can be deformed to canonical form $\tilde{\Psi}^5_0(z) = [1 + z, (2+z)z^4, (1+2z)z, -(4+z)z^3]$.*

Proof. First step is in showing that there exists deformation of $\tilde{\Psi}^5_1(z)$ to $\tilde{\Psi}^5_2(z)$ from theorem 7.8. We will do it by deformation both of $\tilde{\Psi}^5_1(z)$ and $\tilde{\Psi}^5_2(z)$ to $\tilde{\Psi}^5_0(z)$. These deformations are given by $\tilde{\Psi}^5_1(z,t) = [1 + tz, (2t+z)z^4, (t+2z)z, (\frac{-5-3t^2}{2} + tz)z^3]$, $t \in [0,1]$ and $\tilde{\Psi}^5_2(z) = [1+tz, (2t+z)z^4, (1+2tz)z, (-4t+\frac{2t^2-5}{3}z)z^3]$, $t \in [0,1]$ respectively.
Second step is in deforming every curve $\tilde{\Psi}^5(z)$ to one of the two curves $\tilde{\Psi}^5_1(z)$ and $\tilde{\Psi}^5_2(z)$. Here we have three cases

1) $rs = 0$, $lm = -5$ and deformation to $\tilde{\Psi}^5_1(z)$ is given by $\tilde{\Psi}^5_1(z,t) = [1 + a(1-t)z, (h(1-t)+z)z^4, (r(1-t)+lz)z, (m+s(1-t)z)z^3]$, $t \in [0,1]$ and then applying $g_1 = diag\{1, 1, \frac{2}{l}, -\frac{5}{2m}\}$.

2) $lm = 0$, $rs = -\frac{5}{3}$ and deformation to $\tilde{\Psi}^5_2(z)$ is given by $\tilde{\Psi}^5_2(z,t) = [1 + a(1-t)z, (h(1-t)+z)z^4, (r+l(1-t)z)z, (m(1-t)+sz)z^3]$, $t \in [0,1]$ and then applying $g_2 = diag\{1, 1, \frac{1}{r}, -\frac{5}{3s}\}$.

3) $r, l, m, s \neq 0$ and deformation to $\tilde{\Psi}^5_2(z)$ is given by $\tilde{\Psi}^5_2(z,t) = [1 + a(1-t)z, (h(1-t)+z)z^4, (r+l(1-t)z)z, (m(1-t) - \frac{5+(3ah+lm)(1-t)^2}{3r}z)z^3]$, $t \in [0,1]$ and then applying $g_2 = diag\{1, 1, \frac{1}{r}, r\}$.
It is clear from remark 7.10 that degree is preserving throughout all of these deformations. This completes the proof. □

Corollary 7.12. *Spaces $HHol^{LF}_5(\mathbb{C}P^3)$ and $Harm^{\pm}_5(S^4)$ are path-connected.*

Proof. Follows from theorem 7.4, lemma 4.1, lemma 7.11 and path-connectedness and continuity of actions of $PSp(2,\mathbb{C})$ and $PSL(2,\mathbb{C})$. □

Here we end our summary of known results for harmonic maps of S^2 to S^4. For more details one can refer to [6].

Acknowledgments: Author is thankful to Alexei V. Penskoi for helpful remarks and useful conversations.

References

[1] E. Calabi, Minimal immersions of surfaces in euclidean spheres, J. Differential Geometry 1 (1967), 111-125.

[2] E. Calabi, Quelques applications de l'analyse complexe aux surfaces d'aire minima, Topics in Complex Manifolds (Ed. H. Rossi), Les Presses de l'Universit´e de Montr´eal (1968), 59-81.

[3] J. Barbosa, On minimal immersions of S2 into S2m, Trans. Amer. Math. Soc. 210 (1975), 75-106.

[4] R. L. Bryant, Conformal and minimal immersions of compact surfaces into the 4-sphere, J. Differential Geom. 17 (1982), 455–473.

[5] J. Eells, L. Lemaire, A report on harmonic maps, Bull. London Math. Soc. 10, (1978), 1-68.

[6] J. Bolton and L. M. Woodward, Linearly full harmonic 2-spheres in S^4 of area 20π, Internat. J. Math. 12 (2001), 535–554.

[7] J. Bolton and L. M. Woodward, Higher singularities and the twistor fibration $\pi : \mathbb{C}P^3 \to S^4$, Geom.Dedicata 80 (2000), 231–245.

[8] J. Bolton and L. M.Woodward, The space of harmonic two-spheres in the unit four-sphere, Tohoku Math. J. 58 (2006), 231–236

[9] J. Eells, J. H. Sampson, Harmonic mappings of Riemannian manifolds, Amer. J. Math. 86, (1964), 109-160.

[10] Yung-Chow Wong and Yik-Hoi Au-Yeung, An Elementary and Simple Proof of the Connectedness of the Classical Groups, The American Mathematical Monthly Vol. 74, No. 8 (Oct., 1967), pp. 964-966

[11] H. Reckziegel, Horizontal lifts of isometric immersions into the bundle space of a pseudo-Riemannian submersion, in Global Differential Geometry and Global Analysis, Lecture Notes in Mathematics 1156, Springer, 1985, 18-27.

Faculty of Mathematics, National Research University Higher School of Economics, 6 Usacheva Str., 119048, Moscow, Russia

and

Independent University of Moscow, Bolshoy Vlasyevskiy Pereulok 11, 119002, Moscow, Russia

E-mail adress: ravil.gabdurakhmanov@gmail.com